

Winter 1-2021

## 2021 Update Mtg: Spring Frost and Optimal N Fertilizer Rates in 2nd Gen Hybrids

Peter Jeranyama  
*UMass Cranberry Station*, [peterj@umass.edu](mailto:peterj@umass.edu)

Follow this and additional works at: [https://scholarworks.umass.edu/cranberry\\_extension](https://scholarworks.umass.edu/cranberry_extension)



Part of the [Agriculture Commons](#)

---

### Recommended Citation

Jeranyama, Peter, "2021 Update Mtg: Spring Frost and Optimal N Fertilizer Rates in 2nd Gen Hybrids" (2021). *Cranberry Station Extension meetings*. 336.

Retrieved from [https://scholarworks.umass.edu/cranberry\\_extension/336](https://scholarworks.umass.edu/cranberry_extension/336)


This Article is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Station Extension meetings by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

# Spring Frost and Optimal N Fertilizer Rates in 2<sup>nd</sup> Gen Hybrids

Peter Jeranyama

January 26, 2021

# Spring frost frequency (67 days, 2020)



# Optimal Nitrogen Fertilizer Rates in 2<sup>nd</sup> Gen Hybrids



# Objectives

1. Develop N rate response curve for 2<sup>nd</sup> Gen hybrids and compare with native and 1<sup>st</sup> Gen hybrids
2. Determine effect of N rate on fruit yield and vegetative dry matter
3. Evaluate the effect of field fruit rot as influenced by fertilizer N rate
4. Determine effect of N rate on fruit quality (titratable soluble acids, firmness and TAcy).

## Optimal Fertilizer N rates for 2<sup>nd</sup> Generation Hybrids

- N rates for natives and 1<sup>st</sup> generation hybrids are known
- No such rates exist for 2<sup>nd</sup> generation hybrids - Mullica Queen, DeMoranville or Crimson Queen
- 7 N rates (sub) x 5 cultivars (main) x 4 replications in a RCBD at State Bog

## Fertilizer N rates and Cultivars

- N rates (**0, 40 = farmer, 50, 75, 100, 150 & 200** pounds of N/acre)
- Cultivars: **Howes** (native); **Steven** (1<sup>st</sup> Gen-Hybrid); 2<sup>nd</sup> Gen-Hybrids (**Crimson Queen, DeMoranville and Mullica Queen**)



Phenology	Fertilizer Requirements	Rates (40 lb N/Ac)
Early leaf production	Moderate N for adequate upright length	13 lb (100 lb of 13-13-13) June 8, 2020
Fruit set	High demand for N	18lb (100 lb of 18-8-18) July 8, 2020
Bud set	Occurs during fruit set so supplemental N is needed	9 lb (50 lb of 18-8-18) Aug 3, 2020



# Effect of 200 lb/Acre N on vines




Crimson Queen




DeMoranville



Mullica Queen




- ❑ >100 lb N/acre results in decreased fruit yield
- ❑ MQ – shows a clear optimum
- ❑ Howes – any rate >40 lb N/acre was a waste




**Fruit rot increased with increase in N rate**

## N Fertilizer effect on Fruit Firmness



- ❑ CQ and DeM – fruit firmness did not decrease much with N rate
- ❑ MQ, Stevens & Howes showed considerable decrease with N rate

## N Fertilizer effect on Brix



**Degree Brix – decreased  
in all cultivars with  
increase in N rate**

Measure of maturity,  
flavor & sweetness

## SUMMARY

- Fruit yield decreased >100 lb N/acre for 2<sup>nd</sup>-Gen hybrids; 50 lb N/ac for natives
- Vegetative biomass increased with N rate.
- Fruit rot increased with N rate
- Fruit firmness, Brix, Titratable acids decreased with N
- Fruit diameter-CQ & DM increased, others declined with N.

# ACKNOWLEDGMENTS

1. Collaborators: Dr. Giverson Mupambi & Dr. Casey Kennedy
2. Orlie Jeranyama
3. Nancy DeMoranville
4. Rayan Jahrling

